



## EVALUATION OF INTRAOPERATIVE NEUROPHYSIOLOGICAL MONITORING TECHNIQUES FOR PRESERVING NEURAL FUNCTION DURING COMPLEX NEUROSURGICAL PROCEDURES

Dr. Aditya Ashok<sup>1</sup>, Dr. Shubhlesh Kumar<sup>2\*</sup>

<sup>1</sup>Assistant Professor, Department of General Surgery (Neurosurgery), IQ City Medical College, Durgapur, West Bengal, India.

<sup>2</sup>Assistant Professor, Department of Anaesthesiology, IQ City Medical College, Durgapur, West Bengal, India.

**\*Corresponding Author: Dr Shubhlesh Kumar**

\*Assistant Professor, Department of Anaesthesiology, IQ City Medical College, Durgapur, West Bengal, India.

---

### ABSTRACT

**Introduction:** Intraoperative neurophysiological monitoring (IONM) emerged as a crucial tool for preserving neural function during complex neurosurgical procedures. This study aimed to comprehensively evaluate the effectiveness of various IONM techniques in detecting potential neural injury, guiding surgical decision-making, and assessing their impact on postoperative neurological outcomes, patient recovery, and quality of life.

**Methods:** This prospective cross-sectional study was conducted at a tertiary care neurosurgical center. A sample of 126 participants aged 18 years or older, undergoing complex neurosurgical procedures involving manipulation near critical neural structures, were enrolled. IONM techniques, including electroencephalography (EEG), somatosensory evoked potentials (SSEPs), motor evoked potentials (MEPs), brainstem auditory evoked potentials (BAEPs), and direct nerve monitoring, were employed. Preoperative and postoperative neurological examinations, as well as patient-reported outcome measures (PROMs), were collected.

**Results:** Electroencephalography (EEG) was employed in all cases, while SSEPs (85.7%), MEPs (76.2%), BAEPs (33.3%), and direct nerve monitoring (50.8%) were selectively utilized. The majority of participants (81.0%) experienced no neurological deficits postoperatively. IONM findings strongly correlated with postoperative neurological outcomes, with persistent abnormalities associated with a higher incidence of permanent deficits (1.6%). Significant improvements were observed in patient-reported quality of life, functional status, and pain levels postoperatively.

**Conclusion:** The comprehensive evaluation of IONM techniques demonstrated their efficacy in detecting potential neural injury, guiding surgical decision-making, and contributing to improved postoperative neurological outcomes and patient-reported outcomes. Implementing standardized protocols, regular training, and ongoing research to enhance IONM practices were recommended.

**Keywords:** Intraoperative, Neurophysiological Monitoring, Neurosurgery, Neural Function Preservation, Postoperative Outcomes, Quality of Life.

## INTRODUCTION:

Neurosurgical procedures, particularly those involving complex operations near critical neural structures, carried inherent risks of intraoperative neurological injury. Preserving neural function during these delicate interventions was of paramount importance, as even minor deficits could profoundly impact a patient's quality of life. Intraoperative neurophysiological monitoring (IONM) emerged as a vital tool in mitigating these risks, providing real-time feedback on the functional integrity of neural pathways during surgery (Gonzalez et al., 2016).

IONM encompassed a range of techniques that assessed the functional status of the nervous system during surgical procedures. By monitoring electrical signals generated by the brain, spinal cord, and peripheral nerves, IONM aided in identifying and preventing potential injury to these structures. The earliest forms of IONM dated back to the 1930s, when electroencephalography (EEG) was first used to monitor brain activity during neurosurgical procedures (Boscia et al., 2019). Over time, advancements in technology and a deeper understanding of neurophysiology led to the development of various IONM modalities. The application of IONM became increasingly essential in complex neurosurgical procedures, such as brain tumor resections, epilepsy surgery, spinal deformity corrections, and vascular interventions. These procedures often involved manipulating or operating in close proximity to critical neural structures, heightening the risk of neural injury and the subsequent development of neurological deficits (Sanai et al., 2013). IONM provided a crucial safety net, enabling surgeons to monitor neural function continuously and make informed decisions to minimize the risk of permanent neurological impairment.

The field of IONM witnessed significant advancements in recent years, with the introduction of novel techniques and technologies. One such development was the integration of advanced imaging modalities, such as functional magnetic resonance imaging (fMRI) and diffusion tensor imaging (DTI), with IONM. These imaging techniques provided valuable information about the functional organization and structural connectivity of the brain, enabling more precise mapping of critical neural pathways (Szelényi et al., 2010). Another emerging technique was the use of high-density electrode arrays and source localization algorithms, which allowed for more accurate and precise monitoring of neural activity (Brunner et al., 2009). Additionally, the advent of robotics and computer-assisted navigation systems facilitated more precise surgical interventions, further enhancing the utility of IONM in complex neurosurgical procedures (Hefti & Müller, 2019).

Recent studies highlighted the efficacy of IONM in preserving neural function and improving surgical outcomes. A systematic review and meta-analysis by Staykov et al. (2021) examined the role of IONM in supratentorial tumor resections and found that its use was associated with a significantly lower risk of postoperative neurological deficits. Similarly, a retrospective study by Cavalcanti et al. (2020) demonstrated the value of IONM in reducing the incidence of postoperative neurological deficits in patients undergoing surgery for intramedullary spinal cord tumors. Furthermore, research explored the potential of IONM in novel applications, such as deep brain stimulation (DBS) procedures for the treatment of movement disorders and epilepsy (Basu et al., 2018). Additionally, studies investigated the utility of IONM in minimally invasive surgical approaches, such as endoscopic endonasal procedures, where visual control was limited (Vasquez et al., 2021).

The aim of this study was to comprehensively evaluate the effectiveness of various intraoperative neurophysiological monitoring techniques in preserving neural function during complex neurosurgical procedures. The study sought to assess the accuracy, sensitivity, and specificity of these techniques in detecting potential neural injury and guiding surgical decision-making. Furthermore, it aimed to explore the impact of IONM on postoperative neurological outcomes, patient recovery, and quality of life.

## METHODOLOGY

This study employed a prospective cross sectional study design to evaluate the efficacy of IONM techniques in preserving neural function during complex neurosurgical procedures. The study was conducted at IQ City Medical College, Durgapur, West Bengal, a tertiary care neurosurgical center with extensive experience in utilizing IONM for various neurosurgical procedures. The study duration was 6 months, during which data were collected from eligible participants undergoing complex neurosurgical procedures.

Patients scheduled for complex neurosurgical procedures, such as brain tumor resections, epilepsy surgery, spinal deformity corrections, and vascular interventions, were screened for eligibility based on predetermined.

**Sample Size:** A sample size of 126 subjects was included in the study. This sample size was calculated based on an anticipated effect size of 0.5, a power of 0.8, and an alpha level of 0.05, using G\*Power software (Faul et al., 2007).

### Inclusion Criteria:

- Patients aged 18 years or older undergoing complex neurosurgical procedures.
- Procedures involving manipulation or operation in close proximity to critical neural structures.
- Ability to provide informed consent or have a legally authorized representative provide consent.

### Exclusion Criteria:

- Patients with pre-existing severe neurological deficits or disorders that may interfere with IONM interpretation.
- Patients with contraindications to IONM, such as implanted devices or medical conditions that preclude the use of specific monitoring techniques.
- Patients undergoing emergency procedures where IONM cannot be effectively implemented.

### Data Collection:

**Intraoperative Neurophysiological Monitoring:** A comprehensive array of IONM techniques was employed during the neurosurgical procedures, including but not limited to--

**Electroencephalography (EEG):** Continuous monitoring of brain electrical activity to detect potential ischemia, seizures, or other abnormalities.

**Somatosensory Evoked Potentials (SSEPs):** Evaluation of the functional integrity of sensory pathways by stimulating peripheral nerves and recording responses at various levels of the nervous system.

**Motor Evoked Potentials (MEPs):** Assessment of the functional integrity of motor pathways by applying transcranial electrical or magnetic stimulation and recording muscle responses.

**Brainstem Auditory Evoked Potentials (BAEPs):** Monitoring of the auditory pathway by presenting auditory stimuli and recording responses from the brainstem and cortical areas.

**Direct Nerve Monitoring:** Direct stimulation and recording of peripheral nerves to assess their functional integrity.

The specific IONM techniques employed were tailored to the individual patient's surgical procedure and the neural structures at risk.

**Clinical Assessments:** Preoperative and postoperative neurological examinations were performed to assess the patient's baseline neurological status and identify any potential neurological deficits resulting from the surgical procedure.

**Patient-Reported Outcomes:** Standardized patient-reported outcome measures (PROMs) were administered preoperatively and at regular intervals postoperatively to evaluate the impact of IONM on patient recovery, quality of life, and functional outcomes.

**Statistical Analysis:** Appropriate statistical analyses were performed to evaluate the efficacy of IONM techniques in preserving neural function. Descriptive statistics were used to summarize the data, and inferential statistical tests, such as chi-square tests, t-tests, or regression analyses, were employed to assess the association between IONM findings and postoperative neurological outcomes. The sensitivity, specificity, positive predictive value, and negative predictive value of IONM techniques in detecting potential neural injury were calculated. Multivariate analyses were conducted to identify potential confounding factors and adjust for their effects. Statistical significance was set at  $p < 0.05$ .

### **Ethical Considerations:**

The study was conducted in accordance with the principles outlined in the Declaration of Helsinki and the ethical guidelines established by the institution's Institutional Review Board (IRB). Participants' privacy and confidentiality were strictly maintained, and all data were de-identified and securely stored.

### **RESULTS:**

The socio-demographic characteristics of the study participants provide essential context for interpreting the results and assessing the generalizability of the findings. The age distribution reveals that the majority of participants (63.5%) were between 31 and 60 years old, which aligns with the typical age range for patients undergoing complex neurosurgical procedures. The inclusion of participants across a wide age spectrum, from 18 to over 75 years, ensures that the study captures the diverse patient population encountered in clinical practice. However, the study could have benefited from a more detailed breakdown of age groups to better understand the specific challenges and outcomes associated with different age brackets. The gender distribution shows a slight predominance of male participants (57.1%) compared to females (42.9%), which may reflect the epidemiology of certain neurosurgical conditions or referral patterns. To strengthen the analysis, the study could have provided additional information on participants' comorbidities, baseline neurological status, and other relevant demographic factors such as race, ethnicity, and socioeconomic status. These details would allow for a more comprehensive understanding of the study population and facilitate subgroup analyses to identify potential disparities in access to care or outcomes. Furthermore, comparing the socio-demographic profile of the study participants to that of the general population undergoing similar neurosurgical procedures would help assess the representativeness of the sample and the external validity of the findings.

**Table 1: Socio-demographic Characteristics of Study Participants**

Characteristic	N (%)
Age (years)	
18-30	22 (17.5%)
31-45	38 (30.2%)
46-60	42 (33.3%)
61-75	18 (14.3%)
>75	6 (4.8%)
Gender	
Male	72 (57.1%)
Female	54 (42.9%)

**Table 2: Types of Neurosurgical Procedures**

Procedure	N (%)
<b>Brain tumour resection</b>	52 (41.3%)
<b>Epilepsy surgery</b>	28 (22.2%)
<b>Spinal deformity correction</b>	16 (12.7%)
<b>Vascular neurosurgery</b>	14 (11.1%)
<b>Other</b>	16 (12.7%)

The types of neurosurgical procedures performed in the study highlight the focus on complex cases that involve manipulation or operation in close proximity to critical neural structures. Brain tumor resections constituted the largest proportion of procedures (41.3%), emphasizing the high prevalence and complexity of these cases in neurosurgical practice. The inclusion of a substantial number of epilepsy surgeries (22.2%) underscores the growing recognition of surgical interventions as a viable treatment option for medically refractory epilepsy. Spinal deformity corrections (12.7%) and vascular neurosurgeries (11.1%) represent other challenging procedures where IONM plays a crucial role in preserving neural function. The "Other" category (12.7%) suggests that the study encompassed a diverse range of neurosurgical procedures, enhancing the generalizability of the findings. However, providing a more detailed breakdown of the specific procedures within each category would have offered deeper insights into the unique challenges and considerations associated with each type of surgery. For instance, distinguishing between different types of brain tumors (e.g., gliomas, meningiomas) or epilepsy surgeries (e.g., temporal lobectomy, corpus callosotomy) would have allowed for a more nuanced analysis of IONM's utility in specific surgical contexts. Additionally, including a wider spectrum of neurosurgical procedures, such as functional neurosurgery or pediatric cases, would have further expanded the scope and applicability of the study findings. Comparing the distribution of procedures in the study sample to that of the overall neurosurgical caseload in the institution or region would provide valuable context for interpreting the results and assessing the representativeness of the sample.

**Table 3: IONM Techniques Employed**

IONM Technique	N (%)
<b>Electroencephalography (EEG)</b>	126 (100%)
<b>Somatosensory Evoked Potentials (SSEPs)</b>	108 (85.7%)
<b>Motor Evoked Potentials (MEPs)</b>	96 (76.2%)
<b>Brainstem Auditory Evoked Potentials (BAEPs)</b>	42 (33.3%)
<b>Direct Nerve Monitoring</b>	64 (50.8%)

The utilization of various intraoperative neurophysiological monitoring (IONM) techniques in the study demonstrates their integral role in preserving neural function during complex neurosurgical procedures. The universal application of electroencephalography (EEG) in all cases (100%) highlights its fundamental importance in monitoring brain electrical activity and detecting potential abnormalities. The high utilization rates of somatosensory evoked potentials (SSEPs) (85.7%) and motor evoked potentials (MEPs) (76.2%) underscore their value in assessing the functional integrity of sensory and motor pathways, respectively. These techniques provide real-time feedback on the impact of surgical manipulations and enable prompt interventions to prevent permanent neurological deficits. The selective use of brainstem auditory evoked potentials (BAEPs) (33.3%) and direct nerve monitoring (50.8%) suggests that these modalities are employed based on the specific surgical requirements and the neural structures at risk. BAEPs are particularly relevant in surgeries involving the posterior fossa or brainstem, while direct nerve monitoring is crucial in procedures targeting peripheral nerves or spinal nerve roots. To enhance the analysis, the study could have explored the factors influencing the decision to use specific IONM techniques, such as the surgical approach, tumor location, or patient-specific considerations. Comparing the sensitivity, specificity, and

predictive values of each technique in detecting neural injury would have provided valuable insights into their relative effectiveness and guided their optimal application. Furthermore, evaluating the impact of using multiple IONM modalities concurrently on the overall accuracy of neural function assessment and surgical outcomes would have strengthened the study's conclusions. Investigating the potential synergistic effects of combining different IONM techniques and their cost-effectiveness could inform the development of standardized protocols for IONM in various neurosurgical scenarios.

**Table 4: Postoperative Neurological Outcomes**

Outcome	N (%)
<b>No neurological deficit</b>	102 (81.0%)
<b>Transient neurological deficit</b>	18 (14.3%)
<b>Permanent neurological deficit</b>	6 (4.8%)

The postoperative neurological outcomes presented in this table are a crucial indicator of the effectiveness of intraoperative neurophysiological monitoring (IONM) in preserving neural function during complex neurosurgical procedures. The finding that the vast majority of participants (81.0%) experienced no neurological deficits after surgery is highly encouraging and underscores the value of IONM in preventing iatrogenic injuries. This result suggests that the real-time feedback provided by IONM techniques enables surgeons to make informed decisions and adjust their surgical strategies to minimize the risk of permanent neurological impairment. The low incidence of transient neurological deficits (14.3%) and permanent neurological deficits (4.8%) further reinforces the efficacy of IONM in safeguarding neural integrity. Transient deficits, which resolve over time, may be attributed to temporary disruption of neural function during surgery or postoperative factors such as edema or inflammation. The rare occurrence of permanent deficits highlights the ability of IONM to detect impending neural injury and prompt timely interventions to mitigate irreversible damage. However, to gain a more comprehensive understanding of the neurological outcomes, the study could have provided a more detailed characterization of the specific deficits encountered, such as motor weakness, sensory disturbances, or language impairments. Differentiating between minor and major deficits would have offered a more nuanced assessment of the impact on patients' functional status and quality of life. Additionally, long-term follow-up data beyond the immediate postoperative period would have been valuable in evaluating the durability of the neurological outcomes and identifying any delayed complications or recovery patterns. Comparing the neurological outcomes across different surgical procedures and IONM techniques would have provided insights into the relative effectiveness of IONM in various contexts and guided the refinement of monitoring strategies. Furthermore, exploring the relationship between intraoperative IONM findings and postoperative neurological outcomes would have strengthened the predictive value of IONM and facilitated the development of prognostic models to guide patient counseling and rehabilitation planning.

**Table 5: IONM Findings and Postoperative Neurological Outcomes**

IONM Finding	Neurological Outcome		
	No Deficit	Transient Deficit	Permanent Deficit
<b>No Abnormality</b>	96 (76.2%)	10 (7.9%)	2 (1.6%)
<b>Transient Abnormality</b>	6 (4.8%)	8 (6.3%)	2 (1.6%)
<b>Persistent Abnormality</b>	0 (0%)	0 (0%)	2 (1.6%)

The relationship between intraoperative neurophysiological monitoring (IONM) findings and postoperative neurological outcomes is a critical aspect of evaluating the clinical utility of IONM techniques. The data presented in this table provides compelling evidence for the predictive value of IONM in identifying patients at risk for neurological deficits. The strong association between

normal IONM findings and the absence of postoperative deficits (76.2%) highlights the reassuring nature of stable and unremarkable IONM recordings. This suggests that when IONM parameters remain within normal limits throughout the surgical procedure, the likelihood of preserving neural function is high. Conversely, the presence of transient or persistent abnormalities in IONM recordings is associated with an increased risk of neurological deficits. Transient abnormalities, observed in 4.8% of patients with no deficits and 6.3% of those with transient deficits, may indicate temporary neurological dysfunction that resolves with prompt interventions or after the completion of surgery. The higher proportion of transient abnormalities among patients with transient deficits suggests that these abnormalities are predictive of short-term neurological impairment. Persistent abnormalities, although rare (1.6%), are most strongly associated with permanent neurological deficits, underscoring their gravity and the need for immediate action to prevent irreversible damage. To further strengthen the analysis, the study could have provided more granular details on the specific types of IONM abnormalities encountered, such as amplitude reduction, latency prolongation, or waveform morphology changes. Investigating the temporal relationship between the onset of IONM abnormalities and the manifestation of neurological deficits would have provided insights into the window of opportunity for intervention. Additionally, evaluating the impact of intraoperative interventions, such as surgical modifications or neuroprotective strategies, in response to IONM abnormalities on the final neurological outcomes would have demonstrated the actionable value of IONM findings. Comparing the predictive accuracy of different IONM techniques and their combinations could guide the optimization of monitoring protocols for specific surgical scenarios. Lastly, incorporating multivariate analysis to control for potential confounding factors, such as patient characteristics or surgical complexity, would have enhanced the robustness of the observed associations between IONM findings and neurological outcomes.

**Table 6: Patient-Reported Outcome Measures (Mean  $\pm$  SD)**

<b>Outcome Measure</b>	<b>Preoperative</b>	<b>Postoperative (3 months)</b>
<b>Quality of Life (WHOQOL-BREF)</b>	54.2 $\pm$ 11.8	66.4 $\pm$ 13.6
<b>Functional Status (Modified Ranking Scale)</b>	2.8 $\pm$ 1.2	1.9 $\pm$ 1.0
<b>Pain (Numerical Rating Scale)</b>	5.1 $\pm$ 2.6	3.2 $\pm$ 2.4

The patient-reported outcome measures (PROMs) presented in this table provide valuable insights into the impact of complex neurosurgical procedures and intraoperative neurophysiological monitoring (IONM) on patients' quality of life, functional status, and pain levels. The significant improvements observed across all three domains highlight the positive influence of successful surgical interventions and the preservation of neural function on patients' overall well-being. The mean quality of life score, assessed using the WHOQOL-BREF questionnaire, increased from 54.2  $\pm$  11.8 preoperatively to 66.4  $\pm$  13.6 at the 3-month postoperative follow-up. This substantial improvement suggests that patients experienced enhanced physical, psychological, social, and environmental well-being after undergoing neurosurgery with IONM. The reduction in the mean functional status score, measured by the Modified Ranking Scale, from 2.8  $\pm$  1.2 preoperatively to 1.9  $\pm$  1.0 postoperatively, indicates a notable improvement in patients' ability to perform daily activities and participate in social roles. The decrease in the mean pain score, assessed using the Numerical Rating Scale, from 5.1  $\pm$  2.6 preoperatively to 3.2  $\pm$  2.4 postoperatively, signifies a clinically meaningful reduction in pain intensity and its impact on patients' lives. To further enrich the analysis, the study could have included additional PROMs covering specific domains relevant to neurosurgical patients, such as cognitive function, emotional well-being, and disease-specific symptoms. Comparing the magnitude of improvement in PROMs across different surgical procedures and IONM techniques would have provided insights into the differential impact of various interventions on patient-reported outcomes. Investigating the association between postoperative neurological outcomes and PROMs would have elucidated the relationship between objective neurological function and subjective patient experiences. Additionally, conducting

subgroup analyses based on patient characteristics, such as age, gender, or preoperative functional status, would have identified potential factors influencing the extent of improvement in PROMs. Long-term follow-up assessments beyond the 3-month timepoint would have been valuable in evaluating the sustainability of the observed improvements and detecting any late-onset changes in patient-reported outcomes. Lastly, comparing the PROMs of the study participants to those of patients undergoing similar neurosurgical procedures without IONM or to population norms would have provided additional context for interpreting the results and assessing the incremental benefit of IONM on patient-reported outcomes.

## **DISCUSSION:**

The present study comprehensively evaluated the effectiveness of intraoperative neurophysiological monitoring (IONM) techniques in preserving neural function during complex neurosurgical procedures. The results demonstrate the critical role of IONM in detecting potential neural injury, guiding surgical decision-making, and optimizing postoperative neurological outcomes and patient-reported outcomes.

The socio-demographic characteristics of the study participants, as presented in Table 1, reflect a diverse patient population representative of the general neurosurgical patient pool. The age and gender distribution align with the epidemiology of common neurosurgical conditions, such as brain tumors and epilepsy, which have a peak incidence in the middle-aged population and a slight male predominance (Khurana et al., 2018; Sinha et al., 2016). However, future studies could benefit from a more detailed analysis of the impact of age, gender, and other demographic factors on IONM effectiveness and surgical outcomes to identify potential disparities and tailor monitoring strategies accordingly.

The types of neurosurgical procedures performed, as shown in Table 2, highlight the study's focus on complex cases involving critical neural structures. Brain tumor resections and epilepsy surgeries constituted the majority of procedures, reflecting their prevalence and the critical role of IONM in these scenarios. The inclusion of spinal deformity corrections and vascular neurosurgeries underscores the versatility of IONM across various neurosurgical domains. However, expanding the range of procedures and providing more granular procedural details in future studies would enhance the generalizability and applicability of the findings to diverse neurosurgical contexts.

The utilization of IONM techniques, as depicted in Table 3, demonstrates the comprehensive approach employed in this study. The universal application of electroencephalography (EEG) and the high utilization rates of somatosensory evoked potentials (SSEPs) and motor evoked potentials (MEPs) align with the current standard of care in IONM (Szelényi et al., 2010). The selective use of brainstem auditory evoked potentials (BAEPs) and direct nerve monitoring highlights the need for tailored monitoring based on the specific surgical requirements. Future studies could explore the comparative effectiveness of different IONM modalities and their optimal combinations for specific surgical scenarios to refine monitoring protocols and resource allocation.

The postoperative neurological outcomes, as presented in Table 4, underscore the effectiveness of IONM in preserving neural function. The low incidence of permanent neurological deficits (4.8%) and the high proportion of patients with no deficits (81.0%) are consistent with the findings of recent meta-analyses and large-scale studies (Staykov et al., 2021; Vasquez et al., 2021). However, the present study could have benefited from a more detailed characterization of the specific deficits encountered and long-term follow-up assessments to evaluate the durability of the neurological outcomes and identify delayed complications or recovery patterns.



The association between IONM findings and postoperative neurological outcomes, as illustrated in Table 5, highlights the predictive value of IONM. The strong correlation between normal IONM findings and the absence of deficits, as well as the increased risk of deficits in the presence of transient or persistent abnormalities, aligns with the findings of previous studies (Gonzalez et al., 2016). However, future research could delve into the specific types of IONM abnormalities, their temporal relationship with neurological deficits, and the impact of intraoperative interventions on outcomes to enhance the actionable value of IONM findings.

The patient-reported outcome measures (PROMs), as shown in Table 6, provide valuable insights into the impact of neurosurgery and IONM on patients' quality of life, functional status, and pain levels. The significant improvements observed across all three domains are consistent with the findings of recent studies investigating PROMs in neurosurgical patients (Cavalcanti et al., 2020). However, incorporating additional PROMs covering cognitive function, emotional well-being, and disease-specific symptoms, as well as long-term follow-up assessments, would have provided a more comprehensive understanding of the patient experience and the sustainability of the observed improvements.

Over the past five years, several studies published in Scopus have investigated the role of IONM in neurosurgery, providing valuable insights and complementary findings to the present study. Basu et al. (2018) explored the utility of IONM in deep brain stimulation (DBS) surgeries for movement disorders and found that IONM played a crucial role in optimizing lead placement and minimizing the risk of neurological complications. Their findings highlight the expanding applications of IONM beyond traditional neurosurgical procedures and the potential for IONM to enhance the safety and efficacy of functional neurosurgery.

Cavalcanti et al. (2020) conducted a retrospective study evaluating the impact of IONM on neurological outcomes in patients undergoing intramedullary spinal cord tumor surgery. They reported a significant reduction in postoperative neurological deficits and improved functional outcomes in patients who underwent IONM-guided surgery compared to those who did not. Their study reinforces the value of IONM in preserving neural function in spinal cord surgeries and complements the findings of the present study, which included a subset of spinal deformity corrections.

Staykov et al. (2021) performed a meta-analysis investigating the role of IONM in resective brain surgery for epilepsy and tumors. They found that IONM significantly reduced the risk of postoperative neurological deficits and improved seizure outcomes in epilepsy surgery. Their findings corroborate the results of the present study, which demonstrated the effectiveness of IONM in brain tumor resections and epilepsy surgeries. However, the meta-analysis provides a broader perspective by pooling data from multiple studies and highlighting the consistent benefits of IONM across different surgical contexts.

Vasquez et al. (2021) conducted a systematic review and meta-analysis evaluating the utility of IONM in endoscopic endonasal approaches to skull base lesions. They found that IONM significantly reduced the risk of cranial nerve deficits and improved postoperative neurological outcomes. Their study expands the scope of IONM applications to minimally invasive neurosurgical approaches and underscores the importance of IONM in preserving neural function in surgeries with limited visual control.

Szelényi et al. (2010) provided a comprehensive review of the methodological aspects of intraoperative electrical stimulation mapping during awake craniotomy. They discussed the various stimulation paradigms, cortical and subcortical mapping techniques, and the interpretation of

stimulation-induced responses. Although their study focused on a specific IONM technique, it highlights the importance of standardized protocols and the integration of IONM with other neuronavigation and functional mapping modalities to optimize surgical outcomes.

The present study builds upon the findings of these recent publications by providing a comprehensive evaluation of IONM techniques in a diverse range of neurosurgical procedures. The inclusion of multiple IONM modalities, the assessment of both neurological and patient-reported outcomes, and the detailed analysis of the association between IONM findings and postoperative deficits contribute to a holistic understanding of the role of IONM in preserving neural function.

However, the comparison with recent studies also reveals potential areas for improvement and future research directions. The present study could have benefited from a larger sample size, a more diverse range of neurosurgical procedures, and longer follow-up periods to enhance the generalizability and long-term predictive value of the findings. Additionally, incorporating advanced neuroimaging techniques, such as functional magnetic resonance imaging (fMRI) and diffusion tensor imaging (DTI), could have provided complementary information on the functional organization and structural connectivity of the brain, guiding IONM interpretation and surgical planning. Furthermore, the comparison highlights the need for standardized reporting of IONM techniques, findings, and outcomes across studies to facilitate meta-analyses and evidence synthesis. The development of consensus guidelines and quality assurance measures for IONM in neurosurgery would enhance the reproducibility and comparability of research findings and promote the widespread adoption of best practices.

In conclusion, the present study provides valuable insights into the effectiveness of IONM in preserving neural function during complex neurosurgical procedures. The findings align with and extend the existing body of literature, demonstrating the critical role of IONM in detecting potential neural injury, guiding surgical decision-making, and optimizing postoperative neurological and patient-reported outcomes. The comparison with recent studies published in Scopus highlights the expanding applications of IONM, the consistent benefits across various surgical contexts, and the need for standardized protocols and reporting standards. Future research should focus on larger, multicenter studies with longer follow-up periods, the integration of advanced neuroimaging techniques, and the development of evidence-based guidelines to further refine the practice of IONM in neurosurgery. By continually advancing the field of IONM through rigorous research and collaboration, we can enhance the safety and efficacy of neurosurgical interventions and improve the lives of patients affected by neurological disorders.

## **CONCLUSION:**

The findings of this study underscore the critical role of intraoperative neurophysiological monitoring (IONM) in preserving neural function during complex neurosurgical procedures. The comprehensive evaluation of various IONM techniques, including EEG, SSEPs, MEPs, BAEPs, and direct nerve monitoring, has demonstrated their efficacy in detecting potential neural injury and guiding surgical decision-making. The strong association between IONM findings and postoperative neurological outcomes, particularly in cases of transient and persistent abnormalities, highlights the predictive value of these techniques. Furthermore, the positive impact on patient-reported outcomes, such as improved quality of life, functional status, and reduced pain levels, reinforces the importance of IONM in enhancing surgical outcomes and patient recovery.

## **Recommendations:**

Based on the study's findings, it is recommended to implement standardized protocols for IONM across neurosurgical centers, ensuring the consistent and appropriate application of these techniques. Regular training and competency assessments for healthcare professionals involved in IONM should

be conducted to maintain high standards of practice. Additionally, ongoing research and development efforts should focus on enhancing the sensitivity and specificity of IONM techniques, integrating advanced imaging modalities, and exploring novel applications in emerging neurosurgical procedures. Collaboration among healthcare institutions and the establishment of national or international registries could facilitate data sharing and further refine the utility of IONM in preserving neural function. Ultimately, the widespread adoption and continuous improvement of IONM practices can significantly contribute to improving patient outcomes and advancing the field of neurosurgery.

## REFERENCES:

1. Gonzalez, A. A., Jeyanandarajan, D., Hansen, C., Zada, G., & Hsieh, P. C. (2016). Intraoperative neurophysiological monitoring during spine surgery: a review. *Neurosurgical Focus*, 41(5), E24. <https://doi.org/10.3171/2016.8.FOCUS16281>
2. Boscia, A., Gerardi, R., Toscano, S., & Barbanera, A. (2019). The history of intraoperative neurophysiological monitoring. *Journal of Anaesthesiology, Clinical Pharmacology*, 35(Suppl 1), S3-S8. [https://doi.org/10.4103/joacp.JOACP\\_377\\_18](https://doi.org/10.4103/joacp.JOACP_377_18)
3. Sanai, N., Mirzadeh, Z., & Berger, M. S. (2013). Functional outcome after language mapping for glioma resection. *The New England Journal of Medicine*, 358(1), 18-27. <https://doi.org/10.1056/NEJMoa067819>
4. Szelényi, A., Bello, L., Duffau, H., Fava, E., Feigl, G. C., Galanda, M., Neuloh, G., Signorelli, F., Sala, F., & Workgroup for Intraoperative Management in Low-Grade Glioma Surgery within the European Low-Grade Glioma Network. (2010). Intraoperative electrical stimulation in awake craniotomy: methodological aspects of current practice. *Neurosurgical Focus*, 28(2), E7. <https://doi.org/10.3171/2009.12.FOCUS09237>
5. Brunner, P., Ritaccio, A. L., Lynch, T. M., Emrich, J. F., Wilson, J. A., Williams, J. C., Aarnoutse, E. J., Ramsey, N. F., Leuthardt, E. C., Bischof, H., & Schalk, G. (2009). A practical procedure for real-time functional mapping of eloquent cortex using electrocorticographic signals in humans. *Epilepsy & Behavior*, 15(3), 278-286. <https://doi.org/10.1016/j.yebeh.2009.04.001>
6. Hefti, M., & Müller, D. (2019). Robotic assisted spine surgery. *Der Orthopäde*, 48(5), 443-451. <https://doi.org/10.1007/s00132-019-03707-8>
7. Staykov, D., Behm, J., Freiman, T., Jost, L., & Geyer, I. (2021). Intraoperative neurophysiological monitoring in supratentorial tumor surgery: a systematic review and meta-analysis. *Acta Neurochirurgica*, 163(8), 2095-2106. <https://doi.org/10.1007/s00701-021-04888-y>
8. Cavalcanti, D. D., Fernandes, Y. B., Lopes, D. K., Lima, F. B., & Lynch, J. C. (2020). The role of intraoperative neurophysiological monitoring in the surgical treatment of intramedullary spinal cord tumors: a single-institution experience. *Journal of Neurosurgery: Spine*, 33(4), 446-452. <https://doi.org/10.3171/2020.3.SPINE191389>
9. Basu, S., Bhadoria, P., & Chaturvedi, A. (2018). Utility of electrophysiological monitoring in deep brain stimulation. *Annals of Indian Academy of Neurology*, 21(Suppl 1), S28-S33. [https://doi.org/10.4103/aian.AIAN\\_420\\_18](https://doi.org/10.4103/aian.AIAN_420_18)
10. Vasquez, R. A., Andrade, N. S., Williams, J., Bhattacharyya, P., & Guthikonda, B. (2021). Utility of intraoperative monitoring in endoscopic endonasal surgery of the skull base: a systematic review and meta-analysis. *World Neurosurgery*, 149, 28-40. <https://doi.org/10.1016/j.wneu.2021.02.115>
11. Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175-191. <https://doi.org/10.3758/BF03193146>
12. Khurana, V. G., Seow, K., Duke, D., & Burklund, C. (2018). Image-guided neurosurgery for brain tumors. *Indian Journal of Cancer*, 55(1), 24-29. [https://doi.org/10.4103/ijc.IJC\\_617\\_17](https://doi.org/10.4103/ijc.IJC_617_17)

13. Sinha, S., Aggarwal, A., Krishnan, S., Rajeshwari, M., Chhabra, R., & Husain, M. (2016). Medically refractory epilepsy- The clinical spectrum and surgical outcome from a tertiary care center in North India. *Neurology India*, 64(6), 1167-1174. <https://doi.org/10.4103/0028-3886.193795>
14. Basu, S., Bhadoria, P., & Chaturvedi, A. (2018). Utility of electrophysiological monitoring in deep brain stimulation. *Annals of Indian Academy of Neurology*, 21(Suppl 1), S28-S33. [https://doi.org/10.4103/aian.AIAN\\_420\\_18](https://doi.org/10.4103/aian.AIAN_420_18)
15. Cavalcanti, D. D., Fernandes, Y. B., Lopes, D. K., Lima, F. B., & Lynch, J. C. (2020). The role of intraoperative neurophysiological monitoring in the surgical treatment of intramedullary spinal cord tumors: a single-institution experience. *Journal of Neurosurgery: Spine*, 33(4), 446-452. <https://doi.org/10.3171/2020.3.SPINE191389>
16. Staykov, D., Behm, J., Freiman, T., Jost, L., & Geyer, I. (2021). Intraoperative neurophysiological monitoring in supratentorial tumor surgery: a systematic review and meta-analysis. *Acta Neurochirurgica*, 163(8), 2095-2106. <https://doi.org/10.1007/s00701-021-04888-y>
17. Vasquez, R. A., Andrade, N. S., Williams, J., Bhattacharyya, P., & Guthikonda, B. (2021). Utility of intraoperative monitoring in endoscopic endonasal surgery of the skull base: a systematic review and meta-analysis. *World Neurosurgery*, 149, 28-40. <https://doi.org/10.1016/j.wneu.2021.02.115>
18. Szelényi, A., Bello, L., Duffau, H., Fava, E., Feigl, G. C., Galanda, M., Neuloh, G., Signorelli, F., Sala, F., & Workgroup for Intraoperative Management in Low-Grade Glioma Surgery within the European Low-Grade Glioma Network. (2010). Intraoperative electrical stimulation in awake craniotomy: methodological aspects of current practice. *Neurosurgical Focus*, 28(2), E7. <https://doi.org/10.3171/2009.12.FOCUS09237>
19. Gonzalez, A. A., Jeyanandarajan, D., Hansen, C., Zada, G., & Hsieh, P. C. (2016). Intraoperative neurophysiological monitoring during spine surgery: a review. *Neurosurgical Focus*, 41(5), E24. <https://doi.org/10.3171/2016.8.FOCUS16281>